

A Cost Analysis of Treating Pediatric Dental Patients Using General Anesthesia Versus Conscious Sedation

Jessica Y. Lee, DDS, MPH, William F. Vann Jr, DMD, PhD, Michael W. Roberts, DDS

Department of Pediatric Dentistry, University of North Carolina, Chapel Hill, North Carolina
Winner of the 2001 Anesthesia Research Foundation Essay Award.

The purpose of this pilot study is to report a cost comparison of general anesthesia (GA) versus oral conscious sedation (CS) for pediatric dental patients. The study sample included 22 children whose parents or guardians selected GA care for their child. Selection criteria limited inclusion to healthy children (American Society of Anesthesiologists' classification I) aged 24–60 months. The subjects acted as their own comparison group to an estimation CS model. Models were developed to assess societal costs for treatment under GA and CS. Treatment rendered was equalized using the dental relative based value unit scale.

Key Words: Cost analysis; Pediatric dentistry; General anesthesia; Conscious sedation.

An overwhelming majority of pediatric dental patients can be treated in the conventional dental environment without the use of pharmacologic agents, except for the occasional use of nitrous oxide or oxygen inhalation analgesia for the mildly anxious child. Pharmacologic management may be essential to provide invasive dental procedures for children who are developmentally or medically compromised, and sometimes this modality is the treatment of choice for preschool-aged children who have not developed the language skills or attention span to cope with conventional dental care. For such children, the 2 most popular modalities of pharmacologic management are general anesthesia (GA) and conscious sedation (CS).¹

General anesthesia is a controlled state of unconsciousness that is accompanied by loss of protective reflexes, including the ability to maintain an airway independently and to respond purposefully to physical stimulation or verbal commands.² Conscious sedation is a minimally depressed level of consciousness that does not affect a patient's abilities to maintain a patent airway

independently and continuously or to respond appropriately to physical stimulation or verbal commands.³

Both GA and CS have higher levels of risks and costs than does conventional care. Although GA is viewed as the more expensive modality of treatment when direct costs only are assessed, this is not clear when societal costs are considered.⁴ Economists generally agree that to calculate societal costs one must also consider opportunity cost. A broad definition of opportunity cost might be as follows: an action that represents the value of next most valuable action forgone or the value of the next best alternative that a decision forces the decision maker to forgo. For example, if a dentist decides to take a vacation from practice, the opportunity cost for this time off is the wage lost for the time out of the office. Gold and colleagues⁵ state that "the best approximation of the opportunity cost for adults is the wage they are, or could be making at work." For the purposes of this study, opportunity cost is defined as income forgone due to dental treatment.⁶ The societal cost perspective is indeed even larger than the parents' or family's perspective. For example, income forgone for parents' wages is relatively easy to calculate, but there is also a societal cost for children missing school due to pain or dental appointments or for school nurses or teachers who must take time with such children. Although such

Received July 1, 2001; accepted for publication July 11, 2001.

Address correspondence to Jessica Y. Lee, DDS, MPH, Department of Pediatric Dentistry, University of North Carolina, 220 Brauer Hall, CB 7450, Chapel Hill, NC 27599-7450.

Anesth Prog 48:82–88 2001

© 2001 by the American Dental Society of Anesthesiology

ISSN 0003-3006/01/\$9.50
SSDI 0003-3006(01)

societal costs are incalculable for the purposes of this study, the authors offer this as a part of the rationale for the importance of taking the societal perspective into account.

Little has been reported in the literature relative to societal cost comparisons for children's dental care. A recent study compared the use of inhalation sedation via nitrous oxide or oxygen versus outpatient general anesthesia for extractions and minor oral surgery in pediatric dental patients.⁴ The patients were scheduled for 1 sedation appointment and 1 GA appointment. In examining the parents' satisfaction and costs of both treatments, the investigators concluded that for extraction-only treatment, it was more cost-effective to use inhalation sedation than GA. The costs in the study were based on direct cost only, excluding indirect and opportunity costs.

Many factors can influence a parent's decision to choose GA or CS for their child. Consideration might include risks and safety of the procedure, their child's perceived comfort, the parent's assessment of the child's cooperation, the impact of the procedure on the child's developing psyche, the amount of care needed, the probability that treatment can be completed with the given modality, and the cost of care.^{7,8} Despite the widespread use of both GA and CS, there have been few cost analysis studies of either modality, and there are no reports of cost analyses from the societal perspective for pediatric dental patients.⁹

The specific aims of this study were as follows: (a) to determine the societal costs for treating pediatric dental patients using GA and CS, (b) to determine the relationship between cost and treatment need for GA and CS, and (c) to determine the relationship between GA versus CS cost models.

MATERIALS AND METHODS

The 3 principal types of cost-effectiveness studies include (a) the cost-consequence model, (b) the cost-effectiveness model, and (c) the cost-minimization model. The cost-consequence model analyzes only the outcome of interest, under the assumption that there are no differences in costs. As an example, this model might be applied to a comparison of 2 dental materials where the cost of the materials and their application are relatively the same, but the outcome of interest is the material survival over time. The cost-effectiveness model examines the true cost-to-outcome ratio for the treatment and a comparator. Using a similar material science example, this model would be used in a study in which the restorative materials may differ in cost and survival. The cost-minimization model analyzes the cost differential

between 2 treatments. The underlying assumption in this model is that the outcome of either treatment will be equal or similar. Using the material science example, the cost-minimization model would compare the costs of the restorative materials under the assumption that the materials performed the same over time.

This pilot study used a cost-minimization model for outpatient GA versus oral CS. This investigation relied on each individual patient as the unit of analysis and each patient contributed cost data. The analyses used evaluated the societal costs, defined as the sum total of direct, indirect, and opportunity costs. Cost analyses usually use mathematical, estimation, or simulation modeling.⁹ In this study, the GA model relied on primary data and the CS model was an estimation model.

Sample

The sample included 22 children who required pharmacologic management for dental treatment because of short-term situational anxiety in the conventional dental environment. All were scheduled for treatment using GA because their parent or guardian opted for this modality of care. The patients' ages ranged from 24–60 months at the time of the GA appointment. All were healthy children with no contraindications for routine dental care. All met requirements for American Society of Anesthesiologists class I anesthesia risk.

To maintain consistency and reliability, all patients were treated by the same dental operator in the GA setting. The patient sample served as its own comparison group to an estimation model for CS treatment.

Panel of Experts

Expert judgment and consensus panels involve synthesis approaches used to estimate probabilities, costs, preference weights, and other variables in cost-effectiveness studies.⁵ For this study, a panel of 4 experts was used to determine values in the CS estimation model that could not be obtained from actual data. The panel consisted of 2 experts each in the areas of CS and GA. The experts were selected based on their extensive research and clinical experience. All were board-certified pediatric dentists who each have 20–25 years of clinical practice experience in the specialty.

Relative Based Value Units

The treatment rendered was assessed using the dental relative based value units (RBVUs).¹⁰ The RBVUs are based on the time and difficulty of procedures. The RBVU system has been used in medicine for many years as a way to value medical procedures across disciplines

Table 1. Examples of RBVUs for Selected Dental Procedures*

Dental Procedures	ADA Code	RBVU
Posteroanterior x-ray examination (1 film)	0220	0.5
Bitewing x-ray examination (2 films)	0272	0.6
Prophylaxis	1201	0.75
1 surface amalgam, primary	2110	1.0
1 surface resin, primary	2330	2.0
Stainless steel crown	2930	4.0
Pulpotomy, primary	3220	3.0
Extractions, primary	7110	2.0

* RBVUs indicates relative based value units; ADA, American Dental Association.

and specialties. The dental RBVU system was developed in 1985 to equate dental procedures. Normal distribution and SDs and means were used to develop the scale. The data were analyzed for validity, statistically normalized, and weighted.

The RBVUs (Table 1) are considered to be valid and reliable measures of dental procedures. They are used widely by health insurance organizations such as Aetna, Blue Cross/Blue Shield, Prudential, and Delta Dental, as well as Medicaid agencies in Vermont, Kentucky, and South Dakota, to determine the value of procedures.¹⁰

To equalize treatment for the GA and CS models, RBVUs were calculated from the treatment rendered for the 22 children during their GA appointment. The same RBVU data for the GA appointment were used in the CS estimation model, so dental procedures were equal in both models. To determine the number of CS appointments to equate the same RBVUs rendered under GA, this study relied on the clinical judgment of the same operator who completed the 22 GA cases. This approach allowed us to control for operator speed and judgment in determining the number of CS appointments that a given child might need.

Cost Models

Models used in this study were taken from the societal perspective. This approach relies on the perspective of the decision maker, which in this study was the parent or guardian.⁹ The cost models included costs incurred by the family for treatment and opportunity costs. Gusten and colleagues⁶ define opportunity costs as income forgone for the decision of treatment. The accounting data included both indirect and direct costs.

Data for cost models were collected from hospital and dental school accounting records for the patients. There are many ways to calculate opportunity costs; for the purposes of this study, an aggregate measure was used.

Income data by county were obtained for 1997 fiscal year from the Economic Policy Unit of the North Carolina Department of Commerce. The average wage earnings for each adult's county of residence were used to calculate lost income. In summary, opportunity costs in the model were calculated by multiplying the aggregate family wage earnings by the total hours lost because of the child's dental treatment.

Assumptions Underlying Model Development

This study relied on the cost-minimization model for using GA versus CS. In cost analysis studies, many assumptions must be made to develop a model.⁹ For the GA model, this investigation relied on the panel of experts to estimate the parental time commitment for a preoperative GA visit (4 hours) and for the GA appointment itself (8 hours). For the number of adults accompanying the child for appointment procedures, the actual numbers were used, either 1 adult or 2 adults, depending on who accompanied the child.

For the CS model, the panel of experts estimated that the preappointment physical examination would require 4 hours and that only 1 parent would accompany the child for this appointment. For the CS new patient examination, they reached a consensus that this appointment would require a 2-hour appointment and that 1 parent would accompany the child. Finally, they estimated that 2 adults would accompany their child for their CS appointment and a 4-hour time commitment would be required for each sedation visit.

Cost of GA

The cost of GA is determined by the following formula: $(C_{\text{total}} = C_{\text{screen}} + C_{\text{preop}} + C_{\text{tx}} + C_{1\text{st}\frac{1}{2}\text{hr}} + C_{\text{add}\frac{1}{2}\text{hr}} X_1 + C_{\text{anesth}} + C_{\text{rec}} + C_{\text{pt/family}})$. For the GA model, the cost (\$22) of screening (C_{screen}) was obtained from dental school records. The cost (\$88) for the preoperative medical appointment (C_{preop}) was obtained from the hospital accounting records for the preoperative GA evaluation. The actual dental fees for the procedures rendered under GA were used for the cost of treatment (C_{tx}). This cost was equalized to the CS model using RBVUs. The hospital cost for the GA appointment ($C_{1\text{st}\frac{1}{2}\text{hr}}$ and $C_{\text{add}\frac{1}{2}\text{hr}}$) was obtained from the hospital accounting records generated for each patient. Because hospital fees vary from the first half hour (\$950) to each additional half hour (\$530), 2 separate variables were used. The variable representing the cost of each additional half hour ($C_{\text{add}\frac{1}{2}\text{hr}}$) was multiplied by the time beyond the first half hour (X_1). The cost of the anesthesia (C_{anesth}) was \$145 for the first 30 minutes and \$73 for each additional 30 minutes. The recovery room (C_{rec})

costs were \$110 per hour. The operating room time was obtained from hospital records.

The opportunity cost for each patient was calculated by average income for the county of residence multiplied by the time lost for treatment per adult. Opportunity costs were calculated for both the preoperative and GA appointments. All hospital accounting information was obtained from the Office of Cost Accounting at the University of North Carolina Hospitals.

Cost of CS

The cost of CS is determined by the following formula: $(C_{\text{total}} = C_{\text{screen}} + C_{\text{npe}} + C_{\text{ppe}} + C_{\text{tx}} + C_{\text{appt}} X_2 + C_{\text{appt}} X_3 + C_{\text{pt/family}})$. The cost (\$22) of the screening appointment (C_{screen}) and the cost (\$22) of the CS new patient examination (C_{npe}) were obtained from dental school fees. The estimated cost (\$88) of the physician's physical evaluation (C_{ppe}) for CS examination was obtained from the University of North Carolina ambulatory care unit.

As noted already, the panel of experts estimated the time it took for a CS new patient examination, the physician's preoperative physical examination, and the number of adults accompanying a typical child to preoperative appointment and sedation appointments. This information was necessary to quantify the opportunity cost for each family ($C_{\text{pt/family}}$). To estimate the opportunity cost for each sedation appointment, the average income for the county of residence was multiplied by total time lost due to all phases of treatment for each adult accompanying the child. The CS appointment length was estimated at 60–90 minutes by the consensus panel.

The cost of each sedation appointment (C_{appt}) was defined as the fee charged by the dentist to perform the CS procedures. This fee (\$250) reflects a charge for sedation medications, monitoring equipment, and additional personnel dedicated to monitor and assist in the CS area. The C_{appt} was multiplied by the number of appointments (X_2) estimated for the same treatment to be completed under CS as was completed under GA.

As noted previously, the estimated number of CS appointments needed for each child was made by the same dental operator who completed the actual procedures in the operating room for the GA model. This approach allowed the control for the variable clinic speed. Furthermore, the dental operator who made this judgment had extensive experience in the sedation arena. She made these judgments on estimated CS appointments needed on the basis of the patient's treatment plan, quadrants of care needed, and the patient's body weight. Body weight was used because this vari-

able must be considered for the amount of local anesthetic used in a given visit.

After the number of CS appointments was determined, the probability that the patient would need to return for another appointment because CS was aborted due to patient behavior was then estimated. This abortion rate (X_3) in the Pediatric Sedation Clinic of the Department of Pediatric Dentistry at University of North Carolina at Chapel Hill is 5.6%.¹¹ The cost of actual dental treatment (C_{tx}) was equalized to GA treatment using RBVUs.

Data Analyses

The study's first aim was to establish costs representative of the societal costs for treatment under GA and CS. This was accomplished using a combination of opportunity cost and accounting data that represent direct and indirect costs.

The study's second aim was to establish the relationship between dental treatment rendered and total costs. Ordinary least squares regression analyses¹² were used to examine the association between societal costs of treatment and RBVUs. The outcome measure was the societal costs of treatment, and the major explanatory variable was treatment need as measured by RBVUs. Two regression models were used: Cost for GA = $\beta_0 + \beta_1$ (RBVUs) and Cost for CS = $\alpha_0 + \alpha_1$ (RBVUs). The first equation illustrates the relationship between total costs and RBVUs for GA, whereas the second illustrates the same association for CS.

The study's third aim was to determine the relationship between the GA and CS models. The 2 regression equations were plotted against each other, and the relationship of the predicted regression equations illustrates the association between the GA and CS models. STATA Statistical Software was used for all the data analyses.¹³

RESULTS

The sample included 12 boys (55%) and 10 girls (45%). The mean age of the sample was 40 months (SD, 5.1) with an age distribution as follows: 24–35 months, 32%; 36–47 months, 32%; and 48–60 months, 33%.

Table 2 illustrates the GA results. The mean time spent in the operating room was 2 hours 32 minutes. Most cases (64%) were completed in 3 hours or less. The mean GA charge, excluding the cost for dental procedures and opportunity cost, was \$2326. The mean societal cost was \$2698, a figure that includes opportunity costs but excludes costs for dental procedures.

The mean RBVU (treatment rendered) for GA care

Table 2. General Anesthesia Results*

	No. (%) (N = 22)
Operating room time, h (mean, 2 hours 32 minutes)	
1-2	7 (32)
2-3	7 (32)
3-4	5 (23)
≥4	3 (13)
Total GA charges, \$ (mean, \$2326)†	
1000-1500	2 (9)
1500-2000	10 (45)
2000-2500	3 (14)
2500-3000	7 (32)
Total societal costs, \$ (mean, \$2698)‡	
1000-2000	6 (27)
2000-3000	9 (41)
3000-4000	7 (32)
RBVU, treatment rendered (mean, 53.75 ± 10.5)	
0-35	3 (14)
36-70	13 (59)
71-100	6 (27)

* GA indicates general anesthesia; RBVUs, relative based value units.

† Excluding costs for dental procedures and opportunity costs.

‡ Excluding costs for dental procedures but including opportunity costs.

was 53.75 (SD, 10.5). Table 3 illustrates the results when the same RBVU values were then applied to the CS estimation model. In the estimation model, 23% of the sample required 2 sedation appointments to complete treatment, 41% required 3 sedation appointments, and 36% required 4 or more appointments to complete their care. The mean CS charge, excluding costs for dental procedures and opportunity costs, was \$1363, with the majority (54%) of the cases costing less than \$2000. The mean societal cost was \$2203, a figure that includes opportunity costs but no costs for dental procedures.

Two separate ordinary least squares regression models were executed. In both the GA and CS models, the association between RBVUs and total societal costs was found to be highly significant ($P < .01$) with an adjusted R^2 of 0.77 and 0.63, respectively. The magnitude of effect was also significant, with the coefficients being 24.29 (SD, 4.21) for the GA model and 35.17 (SD, 4.33) for the CS model.

The Figure illustrates the relationship between the predicted regressions lines for the GA and CS models. The intersection of the predicted regression equations illustrates the level of treatment need at which there would be diminishing returns for using CS or the point at which cost savings would favor GA. At a RBVU of

Table 3. Conscious Sedation Results*

	No. (%) (N = 22)
No. of appointments	
2	5 (23)
3	9 (41)
>4	8 (36)
Total CS charges, \$ (mean, \$1363)†	
500-1000	5 (23)
1000-1500	9 (41)
1500-2000	6 (27)
2000-2500	2 (9)
Total societal costs, \$ (mean, \$2203)‡	
1000-2000	12 (54)
2000-3000	5 (23)
3000-4000	5 (23)

* CS indicates conscious sedation.


† Excluding costs for dental procedures and opportunity costs.

‡ Excluding costs for dental procedures, but including opportunity costs

66.5 and a cost of \$2677, the total cost for treatment using CS surpasses that for GA.

DISCUSSION

There are several alternative designs that might be used to derive a cost comparison of GA versus CS. This investigation chose the cost-minimization model using a GA sample that served as its own comparator for a CS estimation model. The advantage of this design was ability to control for social and demographic patient effects in both models. This design also reduced the selection



Relationship between general anesthesia (GA) and conscious sedation (CS) predicted regression lines.

that would be introduced by a parental choice of GA or CS on the basis of family convenience factors. Although such factors are important, this design eliminates bias related to parental choices.

The first aim of this study was to determine the societal costs of treating pediatric dental patients using GA and CS. The opportunity cost values that were used ranged from \$73-\$136 per day, depending on the county of residency. The state average was \$105 per day or \$22,583 per year for the average working adult. The mean total societal costs for treating children using GA versus CS was \$2698 and \$2203, respectively.

The second aim was to determine the relationship between societal costs and treatment rendered for both GA and CS models. The goal with this aim was to develop a model that would explain cost of GA and CS. Using regression analyses for the GA model, this study found that the RBVUs explained more than 70% of the variance in cost. For the CS model, the study found that the RBVUs explained more than 60% of the variance in cost. This can be interpreted to mean that treatment rendered, measured in RBVUs, had a significant effect on the costs of both GA and CS.

The third aim was to determine the relationship between GA and CS cost models. When the GA versus CS regression lines were plotted, the intersection represents the point at which the cost of GA and CS would be equal. The study found that at a RBVU of 66.5 and cost of \$2677, CS cost surpasses GA cost. This critical intersection equates to 3.6 CS appointments.

Outcomes for GA Versus CS

One of the major underlying assumptions in this study was that the treatment outcomes for GA versus CS would be the same. The authors recognize that this is a difficult assumption to make. For example, for CS some practitioners may be faced with making treatment decisions with no or poor-quality radiographs. However, the premise in undertaking a cost-minimization study was based on the assumption of all other things being equal, including outcomes of the treatment.

Decisions Based on Risks

In this clinical environment, when parents consider GA versus CS treatment for their children, they have already made an informed decision that no treatment is not an option with which they are comfortable. Typically, these children have more extensive restorative and surgical needs and many have already received emergency care for pain and swelling. For those parents who consider GA versus CS care, it seems logical that their decision should begin with a comparison of risks for the 2 mo-

dalities. However, such risk data are elusive. With reference to pediatric mortality using GA, 1 : 20,000 is cited in the literature by several authors.¹⁴ In the United States there are no published risk data of GA specific to the dental cases. In England and Wales, the dental GA mortality rate remained constant at 1 : 215,000 between 1970 and 1990. These data included all dental GA cases, both pediatric and adult.⁴

There are no published incidence data for the morbidity and mortality of pediatric CS. Goodman and Moore's classic review in 1983 focused on 14 cases of sedation misadventures, but no incidence data were cited. A tragedy of pediatric CS does surface occasionally in the media. However, since the publication of the Guidelines for Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures in 1985, subsequently revised in 1992 by the American Academy of Pediatrics and in 1996 by the American Academy of Pediatric Dentistry, the authors are aware of no pediatric patient fatalities that have occurred when the original 1985 guidelines have been fully used as the standard of care (American Academy of Pediatric Dentistry, written communication, 1999).

A parental decision to choose GA versus CS is difficult to make on the basis of risks, so parents often must consider other factors, one of which is cost. In selecting cost models, this investigation chose a model that values parents' time away from work for appointment activities.

CONCLUSIONS

Under the conditions of the study, the quantity of the treatment rendered based on RBVUs was highly correlated with the total cost of care from the societal perspective, and if a child needed more than 3 CS appointments, the GA option offered cost savings over the CS treatment option.

REFERENCES

1. Nathan JE. Management of a difficult child: a survey of pediatric dentists use of restraints, sedation and general anesthesia. *J Dent Child*. 1989;3:293-301.
2. American Academy of Pediatrics Committee on Drugs. Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures. *Pediatrics*. 1992;5:1110-1115.
3. American Academy of Pediatric Dentistry. Guidelines for the elective use of pharmacologic conscious sedation and deep sedation in pediatric dental patients. *Pediatr Dent* 1996; 18:30-34.

4. Shaw AJ, Meechan JG, Kilpatrick NM, Welbery RR. The use of inhalation sedation and local anesthesia instead of general anesthesia for extraction and minor oral surgery: a prospective study. *Int J Pediatr Dent*. 1996;6:7-11.
5. Gold MR, Siegal JE, Russell LB, Weinstein MC. *Cost Effectiveness in Health and Medicine*. New York, NY: Oxford University Press; 1996.
6. Gusten AJ, Fletcher TM, Warren LT. *Basic Cost Evaluation*. 2nd ed. St Louis, Mo: Mosby Book Inc; 1997:150-273.
7. Kupietzky A, Blumenstyk A. Comparing the behavior of children treated using general anesthesia with those treated using conscious sedation. *J Dent Child*. 1988;5:122-127.
8. Lawrence SM, McTigue DJ, Wilson S. Parental attitudes toward behavior management techniques used in pediatric dentistry. *Pediatr Dent*. 1991;6:199-203.
9. Squires RH, Morris F, Schluterman S, Drwes B, Galyen L, Kendall B. Efficacy, safety, and cost of intravenous sedation versus general anesthesia in children undergoing endoscopic procedures. *Gastrointest Endosc*. 1995;41:99-104.
10. Relative Value Studies Inc. *Relative Values for Dentists*. Denver, Colo: Relative Values Studies Inc; 1997.
11. Leelataweewud P, Vann WF Jr. Adverse events and outcomes of conscious sedation for pediatric patients. *J Dent Res*. 1999;78:518.
12. Gujarati DM. *Basic Econometrics*. 3rd ed. New York, NY: McGraw-Hill Inc; 1995.
13. STATA Corporation. *STATA Programming and Analysis Manuals*. 5th ed. College Station, Tex: STATA Corp; 1998.
14. Litman RS, Perkins FM, Dawson SC. Parental knowledge and attitudes towards discussing risk of death from anesthesia. *Anesth Analg*. 1993;77:256-260.